



Cardinal Health™

# T.E.D.™ anti-embolism stockings

T.E.D.™ anti-embolism stockings have been clinically proven to reduce the risk of developing deep vein thrombosis (DVT) in physician reviewed, published studies on hospitalized patients<sup>1</sup> and to promote increased blood flow velocity in the legs.<sup>2,3,4</sup>

## Assured efficacy


- Address two components of Virchow's Triad: Stasis<sup>2,3,4</sup> and vessel wall damage<sup>5</sup>
- Large thigh length and knee length sizes designed to fit\* high-risk obese population<sup>6</sup>

## Ease of use

- Fits a wide range of patients
- A defined heel pocket is designed to aid correct placement, while toe opening allows easy inspection of the skin and pulse\*

## Safety by design

- Not manufactured with natural rubber latex
- A clinically proven pressure pattern (see reverse) increases blood flow velocity and reduces venous distension<sup>2,3,4</sup>
- A pressure break at the popliteal vein helps to ensure that blood will continue to flow smoothly through this critical area\*
- Interrupted band and 2-ply gusset prevent a tourniquet effect at the femoral vein




Virchow's Triad

**References:** **1.** 2018, Sachdeva A, Dalton M, Lees T. Graduated compression stockings for prevention of deep vein thrombosis. *Cochrane Database Syst Rev.* 2018;11:Cd001484. **2.** Jamieson R, Calderwood CJ, Greer IA. The effect of graduated compression stockings on blood velocity in the deep venous system of the lower limb in the postnatal period. *BJOG : an international journal of obstetrics and gynaecology.* 2007;114(10):1292-4. **3.** Sigel B, Edelstein AL, Felix WR, Jr., Memhardt CR. Compression of the deep venous system of the lower leg during inactive recumbency. *Archives of surgery (Chicago, Ill : 1960).* 1973;106(1):38-43. **4.** Sigel B., et al. Type of Compression for Reducing Venous Stasis. *Archives of Surgery.* February 1975. Vol 110; 171-175 **5.** Coleridge-Smith PD, et al. Deep Vein Thrombosis: Effect of Graduated Compression Stockings on Distension of the Deep Veins of the Calf. *British Journal of Surgery.* June 1991. Vol 78, No. (6): 724-726. **6.** Yang G, De Staercke C, Hooper WC. The effects of obesity on venous thromboembolism: A review. *Open J Prev Med.* 2012;2(4):499-509. doi:10.4236/ojpm.2012.24069


\* Data on File

# Cardinal Health™ T.E.D.™ anti-embolism stockings

## Clinically-proven pressure pattern



## Fitting recommendations



**Knee length** Measure in standing position if possible

1. Measure calf circumference at greatest portion to determine size.
2. Measure the distance from bend of knee to bottom of heel to determine length.



**Thigh length** Measure in standing position if possible

1. Measure upper thigh circumference at the buttock fold. When thigh circumference measures greater than 25 in., select knee length style.
2. Measure calf circumference at greatest portion to determine size.
3. Measure distance from buttock fold to base of heel to determine length.

### Knee length

1. Calf circumference		2. Length			
		Regular		Long	
Small	<12 in.	<16 in.	<b>7071</b>	>16 in.	<b>7339</b>
Medium	12-15 in.	<17 in.	<b>7115-</b>	>17 in.	<b>7480-</b>
Large	15-17.5 in.	<18 in.	<b>7203-</b>	>18 in.	<b>7594</b>
X-Large	17.5-20 in.	<18 in.	<b>7604</b>	>18 in.	<b>7802</b>
XX-Large	20-23 in.	<18 in.	<b>7470LF</b>	>18 in.	<b>7471LF</b>
XXX-Large	23-26 in.	<18 in.	<b>7472LF</b>	>18 in.	<b>7473LF</b>

Available in white

For beige, black and white T.E.D.™ anti-embolism stockings with closed toe for the recuperating patient, call **800.964.5227** for more information



### Thigh length

1. Thigh circumference	2. Calf circumference	3. Length			
		Short <29 in.	Regular 29-33 in.	Long >33 in.	
<25 in.	Small	<12 in.	<b>3071LF-</b>	<b>3130LF</b>	<b>3222LF</b>
	Medium	12-15 in.	<b>3310LF</b>	<b>3416LF-</b>	<b>3549LF</b>
	Large	15-17.5 in.	<b>3634LF</b>	<b>3728LF-</b>	<b>3856LF</b>
25-32 in.	X-Large	17.5-21.5 in.	<b>3180LF</b>	<b>3181LF</b>	<b>3182LF</b>
32-36 in.	XX-Large	21.5-26 in.	<b>3183LF</b>	<b>3184LF</b>	<b>3185LF</b>

Available in white

### T.E.D.™ stocking precautions

- Proper sizing and application must be assured for optimal benefit of stockings.

© 2020 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, ESSENTIAL TO CARE and T.E.D. are trademarks of Cardinal Health and may be registered in the US and/or in other countries. All other marks are the property of their respective owners. Lit. No. 2MS20-1195736 (06/2020)



Imported & marketed in India by Nuvigen Lifesciences Pvt. Ltd.  
For more information, contact Nuvigen Lifesciences: L5-B, Dharam Complex Village, Rahanal, Bhiwandi, Thane - 421302. Tel: +91 9136966042



cardinalhealth.com